

Calculus CP

Summer Assignment Cover Letter

Hello!

I am very excited that you signed up for Calculus CP for next school year! The summer assignment package is attached to this letter, and it is meant to be completed *gradually* over the course of the summer. I will check the packet during the first week of school, and you will be assessed over its contents. We will not spend copious amounts of class time reviewing these concepts, so please be sure that your skills are solid.

Please text the message @2hc76g to the number 81010. This will allow you to sign up for Remind messages. The Remind system allows you to send me messages also! Please do not hesitate to contact me via Remind or via email ([tess.rivero@bss.k12.ok.us](mailto:tess.rivero@bss.k12.ok.us)) if you have questions.

I look forward to a terrific school year!

Sincerely,  
Mrs. Rivero

P.S. Make sure you have one of the following types of graphing calculators: TI-83/84 or TI-Nspire.

Complex Fractions

When simplifying complex fractions, multiply by a fraction equal to 1 which has a numerator and denominator composed of the common denominator of all the denominators in the complex fraction.

Example:

$$\frac{-7 - \frac{6}{x+1}}{\frac{5}{x+1}} = \frac{-7 - \frac{6}{x+1}}{\frac{5}{x+1}} \cdot \frac{x+1}{x+1} = \frac{-7x - 7 - 6}{5} = \frac{-7x - 13}{5}$$

$$\frac{-2 + \frac{3x}{x-4}}{5 - \frac{1}{x-4}} = \frac{-2 + \frac{3x}{x-4}}{5 - \frac{1}{x-4}} \cdot \frac{x(x-4)}{x(x-4)} = \frac{-2(x-4) + 3x(x)}{5(x)(x-4) - 1(x)} = \frac{-2x + 8 + 3x^2}{5x^2 - 20x - x} = \frac{3x^2 - 2x + 8}{5x^2 - 21x}$$

Simplify each of the following.

1.  $\frac{\frac{25}{a} - a}{5 + a}$

2.  $\frac{2 - \frac{4}{x+2}}{5 + \frac{10}{x+2}}$

3.  $\frac{4 - \frac{12}{2x-3}}{5 + \frac{15}{2x-3}}$

4.  $\frac{\frac{x}{x+1} - \frac{1}{x}}{\frac{x}{x+1} + \frac{1}{x}}$

5.  $\frac{1 - \frac{2x}{3x-4}}{x + \frac{32}{3x-4}}$

## Functions

To evaluate a function for a given value, simply plug the value into the function for  $x$ .

Recall:  $(f \circ g)(x) = f(g(x))$  OR  $f[g(x)]$  read "f of g of x" Means to plug the inside function (in this case  $g(x)$ ) in for  $x$  in the outside function (in this case,  $f(x)$ ).

Example: Given  $f(x) = 2x^2 + 1$  and  $g(x) = x - 4$  find  $f(g(x))$ .

$$\begin{aligned}f(g(x)) &= f(x - 4) \\ &= 2(x - 4)^2 + 1 \\ &= 2(x^2 - 8x + 16) + 1 \\ &= 2x^2 - 16x + 32 + 1 \\ f(g(x)) &= 2x^2 - 16x + 33\end{aligned}$$

Let  $f(x) = 2x + 1$  and  $g(x) = 2x^2 - 1$ . Find each.

6.  $f(2) =$  \_\_\_\_\_      7.  $g(-3) =$  \_\_\_\_\_      8.  $f(t+1) =$  \_\_\_\_\_

9.  $f[g(-2)] =$  \_\_\_\_\_      10.  $g[f(m+2)] =$  \_\_\_\_\_      11.  $\frac{f(x+h) - f(x)}{h} =$  \_\_\_\_\_

Let  $f(x) = \sin x$ . Find each exactly.

12.  $f\left(\frac{\pi}{2}\right) =$  \_\_\_\_\_      13.  $f\left(\frac{2\pi}{3}\right) =$  \_\_\_\_\_

Let  $f(x) = x^2$ ,  $g(x) = 2x + 5$ , and  $h(x) = x^2 - 1$ . Find each.

14.  $h[f(-2)] =$  \_\_\_\_\_      15.  $f[g(x-1)] =$  \_\_\_\_\_      16.  $g[h(x^3)] =$  \_\_\_\_\_

Find  $\frac{f(x+h) - f(x)}{h}$  for the given function  $f$ .

17.  $f(x) = 9x + 3$

18.  $f(x) = 5 - 2x$

### Intercepts and Points of Intersection

To find the x-intercepts, let  $y = 0$  in your equation and solve.  
To find the y-intercepts, let  $x = 0$  in your equation and solve.

**Example:**  $y = x^2 - 2x - 3$

x - int. (Let  $y = 0$ )

$$0 = x^2 - 2x - 3$$

$$0 = (x - 3)(x + 1)$$

$$x = -1 \text{ or } x = 3$$

x - intercepts  $(-1, 0)$  and  $(3, 0)$

y - int. (Let  $x = 0$ )

$$y = 0^2 - 2(0) - 3$$

$$y = -3$$

y - intercept  $(0, -3)$

Find the x and y intercepts for each.

19.  $y = 2x - 5$

20.  $y = x^2 + x - 2$

21.  $y = x\sqrt{16 - x^2}$

22.  $y^2 = x^3 - 4x$

Use substitution or elimination method to solve the system of equations.

Example:

$$x^2 + y - 16x + 39 = 0$$

$$x^2 - y^2 - 9 = 0$$

Elimination Method

$$2x^2 - 16x + 30 = 0$$

$$x^2 - 8x + 15 = 0$$

$$(x - 3)(x - 5) = 0$$

$$x = 3 \text{ and } x = 5$$

Plug  $x = 3$  and  $x = 5$  into one original

$$3^2 - y^2 - 9 = 0 \quad 5^2 - y^2 - 9 = 0$$

$$-y^2 = 0$$

$$16 = y^2$$

$$y = 0$$

$$y = \pm 4$$

Points of Intersection  $(5, 4)$ ,  $(5, -4)$  and  $(3, 0)$

Substitution Method

Solve one equation for one variable.

$$y^2 = -x^2 + 16x - 39 \quad (\text{1st equation solved for } y)$$

$$x^2 - (-x^2 + 16x - 39) - 9 = 0 \quad \text{Plug what } y^2 \text{ is equal to into second equation.}$$

$$2x^2 - 16x + 30 = 0$$

*(The rest is the same as previous example)*

$$x^2 - 8x + 15 = 0$$

$$(x - 3)(x - 5) = 0$$

$$x = 3 \text{ or } x = 5$$

Find the point(s) of intersection of the graphs for the given equations.


23.  $x + y = 8$   
 $4x - y = 7$

24.  $x^2 + y = 6$   
 $x + y = 4$

25.  $x^2 - 4y^2 - 20x - 64y - 172 = 0$   
 $16x^2 + 4y^2 - 320x + 64y + 1600 = 0$

Interval Notation

26. Complete the table with the appropriate notation or graph.

Solution	Interval Notation	Graph
$-2 < x \leq 4$		
	$[-1, 7)$	
		

Solve each equation. State your answer in BOTH interval notation and graphically.

27.  $2x - 1 \geq 0$

28.  $-4 \leq 2x - 3 < 4$

29.  $\frac{x}{2} - \frac{x}{3} > 5$

### Domain and Range

Find the domain and range of each function. Write your answer in INTERVAL notation.

30.  $f(x) = x^2 - 5$

31.  $f(x) = -\sqrt{x+3}$

32.  $f(x) = 3\sin x$

33.  $f(x) = \frac{2}{x-1}$

### Inverses

To find the inverse of a function, simply switch the x and the y and solve for the new "y" value.

Example:

$f(x) = \sqrt[3]{x+1}$	Rewrite f(x) as y
$y = \sqrt[3]{x+1}$	Switch x and y
$x = \sqrt[3]{y+1}$	Solve for your new y
$(x)^3 = (\sqrt[3]{y+1})^3$	Cube both sides
$x^3 = y+1$	Simplify
$y = x^3 - 1$	Solve for y
$f^{-1}(x) = x^3 - 1$	Rewrite in inverse notation

Find the inverse for each function.

34.  $f(x) = 2x + 1$

35.  $f(x) = \frac{x^2}{3}$

Also, recall that to PROVE one function is an inverse of another function, you need to show that  $f(g(x)) = g(f(x)) = x$

Example:

If:  $f(x) = \frac{x-9}{4}$  and  $g(x) = 4x+9$  show  $f(x)$  and  $g(x)$  are inverses of each other.

$$\begin{aligned} f(g(x)) &= 4\left(\frac{x-9}{4}\right) + 9 \\ &= x - 9 + 9 \\ &= x \end{aligned}$$

$$\begin{aligned} g(f(x)) &= \frac{(4x+9)-9}{4} \\ &= \frac{4x+9-9}{4} \\ &= \frac{4x}{4} \\ &= x \end{aligned}$$

$f(g(x)) = g(f(x)) = x$  therefore they are inverses of each other.

Prove  $f$  and  $g$  are inverses of each other.

36.  $f(x) = \frac{x^3}{2}$      $g(x) = \sqrt[3]{2x}$

37.  $f(x) = 9 - x^2, x \geq 0$      $g(x) = \sqrt{9-x}$

Equation of a line

**Slope intercept form:**  $y = mx + b$

**Vertical line:**  $x = c$  (slope is undefined)

**Point-slope form:**  $y - y_1 = m(x - x_1)$

**Horizontal line:**  $y = c$  (slope is 0)

38. Use slope-intercept form to find the equation of the line having a slope of 3 and a y-intercept of 5.
39. Determine the equation of a line passing through the point (5, -3) with an undefined slope.
40. Determine the equation of a line passing through the point (-4, 2) with a slope of 0.
41. Use point-slope form to find the equation of the line passing through the point (0, 5) with a slope of  $\frac{2}{3}$ .
42. Find the equation of a line passing through the point (2, 8) and parallel to the line  $y = \frac{5}{6}x - 1$ .
43. Find the equation of a line perpendicular to the y-axis passing through the point (4, 7).
44. Find the equation of a line passing through the points (-3, 6) and (1, 2).
45. Find the equation of a line with an x-intercept (2, 0) and a y-intercept (0, 3).



## Radian and Degree Measure

Use  $\frac{180^\circ}{\pi \text{ radians}}$  to get rid of radians and convert to degrees.

Use  $\frac{\pi \text{ radians}}{180^\circ}$  to get rid of degrees and convert to radians.

46. Convert to degrees:      a.  $\frac{5\pi}{6}$                       b.  $\frac{4\pi}{5}$                       c. 2.63 radians

47. Convert to radians:      a.  $45^\circ$                       b.  $-17^\circ$                       c.  $237^\circ$

## Angles in Standard Position

48. Sketch the angle in standard position.

a.  $\frac{11\pi}{6}$                       b.  $230^\circ$                       c.  $-\frac{5\pi}{3}$                       d. 1.8 radians

## Reference Triangles

49. Sketch the angle in standard position. Draw the reference triangle and label the sides, if possible.

a.  $\frac{2}{3}\pi$

b.  $225^\circ$

c.  $-\frac{\pi}{4}$


d.  $30^\circ$

## Unit Circle

You can determine the sine or cosine of a quadrantal angle by using the unit circle. The x-coordinate of the circle is the cosine and the y-coordinate is the sine of the angle.

**Example:**  $\sin 90^\circ = 1$

$$\cos \frac{\pi}{2} = 0$$



50. a.)  $\sin 180^\circ$


b.)  $\cos 270^\circ$

c.)  $\sin(-90^\circ)$



d.)  $\sin \pi$

e.)  $\cos 360^\circ$

f.)  $\cos(-\pi)$



## Graphing Trig Functions



$y = \sin x$  and  $y = \cos x$  have a period of  $2\pi$  and an amplitude of 1. Use the parent graphs above to help you sketch a graph of the functions below. For  $f(x) = A \sin(Bx + C) + K$ ,  $A$  = amplitude,  $\frac{2\pi}{B}$  = period,  $\frac{C}{B}$  = phase shift (positive  $C/B$  shift left, negative  $C/B$  shift right) and  $K$  = vertical shift.

Graph two complete periods of the function.

51.  $f(x) = 5 \sin x$

52.  $f(x) = \sin 2x$

53.  $f(x) = -\cos\left(x - \frac{\pi}{4}\right)$

54.  $f(x) = \cos x - 3$

## Trigonometric Equations:

Solve each of the equations for  $0 \leq x < 2\pi$ . Isolate the variable, sketch a reference triangle, find all the solutions within the given domain,  $0 \leq x < 2\pi$ . Remember to double the domain when solving for a double angle. Use trig identities, if needed, to rewrite the trig functions. (See formula sheet at the end of the packet.)

55.  $\sin x = -\frac{1}{2}$

56.  $2 \cos x = \sqrt{3}$

$$57. \cos 2x = \frac{1}{\sqrt{2}}$$

$$58. \sin^2 x = \frac{1}{2}$$

$$59. \sin 2x = -\frac{\sqrt{3}}{2}$$

$$60. 2\cos^2 x - 1 - \cos x = 0$$

$$61. 4\cos^2 x - 3 = 0$$

$$62. \sin^2 x + \cos 2x - \cos x = 0$$


## Inverse Trigonometric Functions:

Recall: Inverse Trig Functions can be written in one of ways:

$$\arcsin(x)$$

$$\sin^{-1}(x)$$

Inverse trig functions are defined only in the quadrants as indicated below due to their restricted domains.




### Example:

Express the value of "y" in radians.

$$y = \arctan \frac{-1}{\sqrt{3}}$$

Draw a reference triangle.



This means the reference angle is  $30^\circ$  or  $\frac{\pi}{6}$ . So,  $y = -\frac{\pi}{6}$  so that it falls in the interval from

$$\frac{-\pi}{2} < y < \frac{\pi}{2}$$

$$\text{Answer: } y = -\frac{\pi}{6}$$

For each of the following, express the value for "y" in radians.

76.  $y = \arcsin \frac{-\sqrt{3}}{2}$

77.  $y = \arccos(-1)$

78.  $y = \arctan(-1)$


**Example:** Find the value without a calculator.

$$\cos\left(\arctan\frac{5}{6}\right)$$

Draw the reference triangle in the correct quadrant first.

Find the missing side using Pythagorean Thm.

Find the ratio of the cosine of the reference triangle.



$$\cos\theta = \frac{6}{\sqrt{61}}$$

For each of the following give the value without a calculator.


63.  $\tan\left(\arccos\frac{2}{3}\right)$

64.  $\sec\left(\sin^{-1}\frac{12}{13}\right)$

65.  $\sin\left(\arctan\frac{12}{5}\right)$


66.  $\sin\left(\sin^{-1}\frac{7}{8}\right)$

## Circles and Ellipses



$$r^2 = (x-h)^2 + (y-k)^2$$

$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$$




For a circle centered at the origin, the equation is  $x^2 + y^2 = r^2$ , where  $r$  is the radius of the circle.


For an ellipse centered at the origin, the equation is  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , where  $a$  is the distance from the center to the ellipse along the  $x$ -axis and  $b$  is the distance from the center to the ellipse along the  $y$ -axis. If the larger number is under the  $y^2$  term, the ellipse is elongated along the  $y$ -axis. For our purposes in Calculus, you will not need to locate the foci.

Graph the circles and ellipses below:


67.  $x^2 + y^2 = 16$




68.  $x^2 + y^2 = 5$



69.  $\frac{x^2}{1} + \frac{y^2}{9} = 1$



70.  $\frac{x^2}{16} + \frac{y^2}{4} = 1$



Limits

## 6.6 Practice - Factoring Strategy

Factor each completely.

1)  $24az - 18ah + 60yz - 45yh$

3)  $5u^2 - 9uv + 4v^2$

5)  $-2x^3 + 128y^3$

7)  $5n^3 + 7n^2 - 6n$

9)  $54u^3 - 16$

11)  $n^2 - n$

13)  $x^2 - 4xy + 3y^2$

15)  $9x^2 - 25y^2$

17)  $m^2 - 4n^2$

19)  $36b^2c - 16xd - 24b^2d + 24xc$

21)  $128 + 54x^3$

23)  $2x^3 + 6x^2y - 20y^2x$

25)  $n^3 + 7n^2 + 10n$

27)  $27x^3 - 64$

29)  $5x^2 + 2x$

31)  $3k^3 - 27k^2 + 60k$

33)  $mn - 12x + 3m - 4xn$

35)  $16x^2 - 8xy + y^2$

37)  $27m^2 - 48n^2$

39)  $9x^3 + 21x^2y - 60y^2x$

41)  $2m^2 + 6mn - 20n^2$

2)  $2x^2 - 11x + 15$

4)  $16x^2 + 48xy + 36y^2$

6)  $20uv - 60u^3 - 5xv + 15xu^2$

8)  $2x^3 + 5x^2y + 3y^2x$

10)  $54 - 128x^3$

12)  $5x^2 - 22x - 15$

14)  $45u^2 - 150uv + 125v^2$

16)  $x^3 - 27y^3$

18)  $12ab - 18a + 6nb - 9n$

20)  $3m^3 - 6m^2n - 24n^2m$

22)  $64m^3 + 27n^3$

24)  $3ac + 15ad^2 + x^2c + 5x^2d^2$

26)  $64m^3 - n^3$

28)  $16a^2 - 9b^2$

30)  $2x^2 - 10x + 12$

32)  $32x^2 - 18y^2$

34)  $2k^2 + k - 10$

36)  $v^2 + v$

38)  $x^3 + 4x^2$

40)  $9n^3 - 3n^2$

42)  $2u^2v^2 - 11uv^3 + 15v^4$



Beginning and Intermediate Algebra by Tyler Wallace is licensed under a Creative Commons Attribution 3.0 Unported License. (<http://creativecommons.org/licenses/by/3.0/>)



## Answers - Factoring Strategy

- |                                    |                                      |
|------------------------------------|--------------------------------------|
| 1) $3(2a + 5y)(4z - 3h)$           | 22) $(4m + 3n)(16m^2 - 12mn + 9n^2)$ |
| 2) $(2x - 5)(x - 3)$               | 23) $2x(x + 5y)(x - 2y)$             |
| 3) $(5u - 4v)(u - v)$              | 24) $(3a + x^2)(c + 5d^2)$           |
| 4) $4(2x + 3y)^2$                  | 25) $n(n + 2)(n + 5)$                |
| 5) $2(-x + 4y)(x^2 + 4xy + 16y^2)$ | 26) $(4m - n)(16m^2 + 4mn + n^2)$    |
| 6) $5(4u - x)(v - 3u^2)$           | 27) $(3x - 4)(9x^2 + 12x + 16)$      |
| 7) $n(5n - 3)(n + 2)$              | 28) $(4a + 3b)(4a - 3b)$             |
| 8) $x(2x + 3y)(x + y)$             | 29) $x(5x + 2)$                      |
| 9) $2(3u - 2)(9u^2 + 6u + 4)$      | 30) $2(x - 2)(x - 3)$                |
| 10) $2(3 - 4x)(9 + 12x + 16x^2)$   | 31) $3k(k - 5)(k - 4)$               |
| 11) $n(n - 1)$                     | 32) $2(4x + 3y)(4x - 3y)$            |
| 12) $(5x + 3)(x - 5)$              | 33) $(m - 4x)(n + 3)$                |
| 13) $(x - 3y)(x - y)$              | 34) $(2k + 5)(k - 2)$                |
| 14) $5(3u - 5v)^2$                 | 35) $(4x - y)^2$                     |
| 15) $(3x + 5y)(3x - 5y)$           | 36) $v(v + 1)$                       |
| 16) $(x - 3y)(x^2 + 3xy + 9y^2)$   | 37) $3(3m + 4n)(3m - 4n)$            |
| 17) $(m + 2n)(m - 2n)$             | 38) $x^2(x + 4)$                     |
| 18) $3(2a + n)(2b - 3)$            | 39) $3x(3x - 5y)(x + 4y)$            |
| 19) $4(3b^2 + 2x)(3c - 2d)$        | 40) $3n^2(3n - 1)$                   |
| 20) $3m(m + 2n)(m - 4n)$           | 41) $2(m - 2n)(m + 5n)$              |
| 21) $2(4 + 3x)(16 - 12x + 9x^2)$   | 42) $v^2(2u - 5v)(u - 3v)$           |



Beginning and Intermediate Algebra by Tyler Wallace is licensed under a Creative Commons Attribution 3.0 Unported License. (<http://creativecommons.org/licenses/by/3.0/>)

## Formula Sheet

These should be memorized.

Reciprocal Identities:       $\csc x = \frac{1}{\sin x}$        $\sec x = \frac{1}{\cos x}$        $\cot x = \frac{1}{\tan x}$

Quotient Identities:       $\tan x = \frac{\sin x}{\cos x}$        $\cot x = \frac{\cos x}{\sin x}$

Pythagorean Identities:       $\sin^2 x + \cos^2 x = 1$        $\tan^2 x + 1 = \sec^2 x$        $1 + \cot^2 x = \csc^2 x$

Double Angle Identities:

$\sin 2x = 2 \sin x \cos x$ $\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$	$\cos 2x = \cos^2 x - \sin^2 x$ $= 1 - 2 \sin^2 x$ $= 2 \cos^2 x - 1$
--	---

Logarithms:       $y = \log_a x$  is equivalent to       $x = a^y$

Product property:       $\log_b mn = \log_b m + \log_b n$

Quotient property:       $\log_b \frac{m}{n} = \log_b m - \log_b n$

Power property:       $\log_b m^p = p \log_b m$

Property of equality:      If  $\log_b m = \log_b n$ , then  $m = n$

Change of base formula:       $\log_a n = \frac{\log_b n}{\log_b a}$

Slope-intercept form:       $y = mx + b$

Point-slope form:       $y - y_1 = m(x - x_1)$

Standard form:       $Ax + By + C = 0$

Quadratic Formula:

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Slope:  $\frac{y_2 - y_1}{x_2 - x_1}$

Distance:

$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Area formulas:

Triangle:  $\frac{1}{2}bh$

Circle:  $\pi r^2$

Rectangle:  $lw$

Trapezoid:  $\frac{1}{2}h(b_1 + b_2)$

Volume Formulas:

$V = lwh$  Rectangular Prism

$V = \frac{1}{3}(\text{area of base})h$   
 Pyramid

$V = \pi r^2 h$  cylinder